

taltz<sup>®</sup>  
(ixekizumab)  
injection  
80 mg/mL

# MAKE YOUR MOVE




Try Taltz for as little as \$5 a month\*

\*Terms and Conditions apply. See pages 8 and 9 for details.



## WELCOME TO TALTZ

This guide offers resources, instructions, and support as you begin your journey.



### TABLE OF CONTENTS

SEE WHAT TALTZ CAN DO FOR PSORIATIC ARTHRITIS	4–5	AUTOINJECTOR & PREFILLED SYRINGE	16–17
DISCOVER WHAT TALTZ CAN DO FOR ANKYLOSING SPONDYLITIS	6–7	DOSING FOR PSORIATIC ARTHRITIS	18–19
GETTING TALTZ COVERED	8–9	DOSING FOR ANKYLOSING SPONDYLITIS	20–21
YOUR TALTZ TREATMENT JOURNEY	10–11	HELPFUL INJECTION TIPS/THINGS TO KNOW	22–23
TALTZ TOGETHER™	12	TAKING TALTZ & STORAGE	24–25
WHO'S WHO	13	PURPOSE AND SAFETY SUMMARY	26–27
TALTZ DELIVERY DEVICES	14–15		

Taltz is for adult patients with active ankylosing spondylitis and active psoriatic arthritis. Taltz is also a treatment for adults with moderate to severe plaque psoriasis who may benefit from taking injections or pills (systemic therapy) or phototherapy (treatment using ultraviolet or UV light).

### SELECT IMPORTANT SAFETY INFORMATION

Taltz affects the immune system. It may increase your risk of infections, which can be serious. Do not use Taltz if you have any symptoms of infection, unless your doctor tells you to. If you have a symptom after starting Taltz, call your doctor right away.

Please see the **Safety Summary** on pages 26 and 27. Please see **Instructions for Use** included with your device.

# SEE WHAT TALTZ CAN DO FOR YOU



Psoriatic arthritis can develop slowly with symptoms like joint pain, or it can develop quickly and be severe.

## HOW TALTZ WORKS

Taltz is a biologic that targets interleukin 17A (IL-17A), which is one protein in your body that plays a role in inflammation and mediates bone and cartilage damage. Taltz works to help block IL-17A within your body.\*

\*The relationship between how Taltz works and clinical outcomes has not been determined.

taltz  
(ixekizumab)  
injection  
80 mg/mL



## WHY TALTZ?



### FOR PEOPLE WITH PSORIATIC ARTHRITIS



#### JOINT SYMPTOM IMPROVEMENT

Taltz helped more than half of people experience **joint symptom improvement** at 24 weeks, including **less joint pain and stiffness plus improved mobility**.



#### HELPS STOP PROGRESSION

For some people, Taltz is proven to help **stop the progression of joint damage** caused by psoriatic arthritis at 16 weeks.



#### PERFORM DAILY ACTIVITIES

Some people taking Taltz showed **improvement** at 24 weeks in their **ability to perform daily activities**.



### FOR PEOPLE WITH MODERATE TO SEVERE PLAQUE PSORIASIS



#### 100% CLEAR SKIN<sup>†</sup>

Taltz is the **first and only** treatment of its kind<sup>‡</sup> for moderate to severe plaque psoriasis that offers **a chance at 100% clear skin**.

<sup>†</sup>Up to 90% of people saw significant improvement of their psoriasis plaques and 4 out of 10 achieved 100% clear skin at week 12.

<sup>‡</sup>IL-17A antagonist that includes 100% clear skin in the Prescribing Information.

#### SELECT IMPORTANT SAFETY INFORMATION

Your doctor should check you for tuberculosis (TB) before you start Taltz, and watch you closely for signs of TB during and after treatment with Taltz. If you have TB, or had it in the past, your doctor may treat you for it before you start Taltz.

Please see the **Safety Summary** on pages 26 and 27. Please see **Instructions for Use** included with your device.

# DISCOVER WHAT TALTZ CAN DO FOR ANKYLOSING SPONDYLITIS

**taltz**<sup>®</sup>  
(ixekizumab)  
injection  
80 mg/mL

Ankylosing spondylitis (AS) is a chronic inflammatory disease that primarily affects the spine. If left untreated, it can lead to spinal fusion, reducing mobility.

## HOW TALTZ WORKS

Taltz is a biologic that targets interleukin 17A (IL-17A), which is one protein in your body that plays a role in inflammation and mediates bone and cartilage damage. Taltz works to help block IL-17A within your body.\*

\*The relationship between how Taltz works and clinical outcomes has not been determined.



## WHY TALTZ?

**TALTZ SIGNIFICANTLY IMPROVES THE SYMPTOMS OF AS AT 16 WEEKS**

### **REDUCES INFLAMMATION**

Taltz decreases morning stiffness.

### **IMPROVES BACK PAIN**

Taltz is proven to relieve spinal pain.



### **IMPROVES ABILITY TO PERFORM DAILY ACTIVITIES**

Taltz substantially increases the ability to carry out different types of actions that range from simple to more demanding.



### **REDUCES FATIGUE**

Taltz may improve the tiredness associated with AS.

### **SELECT IMPORTANT SAFETY INFORMATION**

Do not use Taltz if you have had a serious allergic reaction to ixekizumab or any other ingredient in Taltz, such as: swelling of your eyelids, lips, mouth, tongue or throat, trouble breathing, feeling faint, throat or chest tightness, or skin rash. Get emergency help right away if you have any of these reactions. See the Medication Guide that comes with Taltz for a list of ingredients.

Please see the [Safety Summary](#) on pages 26 and 27. Please see [Instructions for Use](#) included with your device.

# GETTING TALTZ COVERED

taltz<sup>®</sup>  
(ixekizumab)  
injection  
80 mg/mL

## THREE WAYS TO GET A SAVINGS CARD

- 1 Check the box for savings card on the Taltz Together™ enrollment form
- 2 Call Taltz Together at 1-844-825-8966
- 3 Online at [taltz.com](https://www.taltz.com)

For commercially insured patients, Taltz Together may help you pay less for your prescription and can help if you're denied coverage or if your insurance company says you need to switch medications.



PAY AS LITTLE AS **\$5**  
OR \$25 A MONTH\*

**\$5**

### TALTZ COVERED BY INSURANCE

If you are commercially insured and your insurance plan covers Taltz, you'll pay as little as \$5 a month for up to 36 months.\*

\*By using the Taltz Savings Card ("Card"), you attest that you meet the eligibility criteria, agree to and will comply with the Terms and Conditions described below: If you meet program eligibility requirements, have commercial drug insurance, and your insurance provider covers Taltz, you are eligible to pay as little as \$5 monthly, subject to a monthly cap of wholesale acquisition cost plus usual and customary pharmacy charges and a separate maximum annual benefit of \$16,000. If you meet program eligibility requirements, have commercial drug insurance, and your insurance provider denies coverage for Taltz, you are eligible to pay \$25 monthly, subject to a monthly and annual cap of wholesale acquisition cost plus usual and customary pharmacy charges. To continue to participate in the \$25 monthly program, patients who qualify are required to initiate a benefits investigation and request for coverage within 6 months post program initiation, as well as 12 months post program initiation, to verify coverage status and potential eligibility for the \$5 monthly program. Once activated, the Card for both the \$5 and \$25 monthly offers is good for up to 36 months from patient qualification into the program and patients must first use their Card by 12/31/2019. Subject to the terms and conditions of the program, patients may be eligible to re-enroll provided they continue to meet program criteria.

Offer void where prohibited by law. Patient is responsible for any applicable taxes, fees, or amounts exceeding monthly or annual caps. **This offer is invalid for patients without commercial insurance coverage or those whose**

**\$25**

### TALTZ NOT COVERED BY INSURANCE

If you are commercially insured but your insurance plan does not cover Taltz, we will work with you and your insurance company to see if they will provide coverage. If you are denied coverage, you will pay as little as \$25 a month for up to 36 months.\*

**prescription claims are eligible to be reimbursed, in whole or in part, by any governmental program, including, without limitation, Medicaid, Medicare, Medicare Part D, Medigap, DOD, VA, TRICARE/CHAMPUS, or any state patient or pharmaceutical assistance program.** If you live in Massachusetts, the Card expires on the earlier of: (i) the expiration date of this Card 12/31/2022; (ii) the date an AB-rated generic equivalent of Taltz becomes available; or (iii) 12/31/2019, absent a change in Massachusetts state law. If you live in California, the Card expires on the earlier of: (i) the expiration date of this Card 12/31/2022 or (ii) the date an FDA-approved therapeutic equivalent of Taltz or over-the-counter product with the same active ingredient becomes available. Available only in the US and Puerto Rico for residents of the US and Puerto Rico.

By accepting this offer, you agree that if you are required to do so under the terms of your insurance coverage for this prescription or are otherwise required to do so by law, you should notify your insurance carrier of your redemption of this Card. This offer cannot be combined or utilized with any other program, discount, discount card, cash discount card, coupon, incentive, or similar offer involving Taltz. It is prohibited for any person to sell, purchase, or trade; or to offer to sell, purchase or trade; or to counterfeit this Card. This offer may be terminated, rescinded, revoked or amended by Lilly USA, LLC, at any time without notice. This Card is not health insurance. Card activation required. This Card expires on 12/31/2022.

# YOUR TALTZ TREATMENT JOURNEY STARTS WITH YOU



The following steps will help you better understand the process of getting started with and staying on Taltz.

taltz  
(ixekizumab)  
injection  
80 mg/mL



## 1 RECEIVE YOUR PRESCRIPTION

If you're commercially insured and have enrolled in Taltz Together, your prescription will be authorized to ship in as little as 2 days.\* Your doctor may then send your prescription to Taltz Together, where someone will be able to help you navigate the insurance approval process.

Prescriptions for Taltz need to be filled by a specialty pharmacy, so sometimes processing your prescription may take a little longer.

## 2 YOU'VE BEEN APPROVED—NOW EXPECT A CALL!

Taltz Together will inform you as to which specialty pharmacies can fill your prescription. The specialty pharmacy then calls you to set up delivery of Taltz to your home. To ensure delivery, please be sure to answer any calls, as it might be your specialty pharmacy.

## 3 TALTZ MEDICATION ARRIVES

Taltz Together can provide additional support, including a free sharps disposal container. Call if you have any questions about starting your Taltz treatment. Please make sure to review the Taltz dosing schedule on pages 18 and 20.

taltz together™  
(ixekizumab)

If you have any other questions, you can call Taltz Together for additional support at **1-844-TALTZ-NOW** (1-844-825-8966). Make sure to save the number for Taltz Together in your contacts.

\*Shipment can be authorized within two business days of submitting a prescription to either a participating specialty pharmacy or Taltz Together and commercial insurance is verified; patient enrolls in the Taltz Savings Card program and schedules shipment. See [Terms and Conditions](#) on pages 8 and 9.



Taltz Together is a support program for those taking Taltz. For guidance throughout your treatment journey, enroll in Taltz Together, where a Companion in Care<sup>™\*</sup> can help you every step of the way.



Once your doctor prescribes Taltz, there are some key players who may reach out to help get you started.



## CHOOSE THE TYPE OF SUPPORT THAT FITS YOUR NEEDS



### INSURANCE SUPPORT

Assistance navigating the insurance approval process to ensure you pay the lowest amount possible.



### INJECTION SUPPORT

Optional in-home or over-the-phone training with a nurse, as well as injection training videos online.



### ONGOING SUPPORT

A Companion in Care will be there when you need them to help you manage your condition and provide educational materials.



### SHARPS DISPOSAL CONTAINER

A free sharps container to dispose of your used Taltz devices. Once the container is full, mail it back free of charge and receive another one.



### TALTZ TOGETHER COMPANION IN CARE

- Coordinates with your doctor's office, your specialty pharmacy, and your insurance, so you don't have to
- Works with your insurance to determine specialty pharmacy options
- Assists you in activating your Taltz Savings Card, if you're eligible



### YOUR SPECIALTY PHARMACY

- Handles biologic medications like Taltz
- Works with other key players to get Taltz approved for you
- Contacts you once your insurance company approves Taltz to set delivery of your medication



### YOUR INSURANCE COMPANY

- Once your insurance receives a prescription from your doctor, they will:
- Determine if Taltz is covered, which specialty pharmacies can fill your prescription, and how much it may cost you

### READY TO ENROLL?

To enroll in Taltz Together, call **1-844-TALTZ-NOW** (1-844-825-8966) or go to [taltz.com](https://www.taltz.com) to sign up now.

\*Please keep in mind that Taltz Together Companions in Care are not medical professionals. Your doctor is your source for medical advice.

# TALTZ DELIVERY DEVICES




Devices designed with you in mind.

taltz<sup>™</sup>  
(ixekizumab)  
injection  
80 mg/mL




## TALTZ IS AVAILABLE IN 2 DIFFERENT DEVICES

### AUTOINJECTOR



### PREFILLED SYRINGE



### EASY TO USE

At first use, over **94% of patients** agreed the Taltz autoinjector was “**easy to use**” and were confident in their ability to use it. Please see Instructions for Use included with the device.

### SELECT IMPORTANT SAFETY INFORMATION

Taltz may cause or worsen symptoms of Crohn’s disease or ulcerative colitis (inflammatory bowel disease). Tell your doctor if you have any of these symptoms or if they get worse: stomach pain, diarrhea, and weight loss.

Please see the [Safety Summary](#) on pages 26 and 27. Please see [Instructions for Use](#) included with your device.

# AUTOINJECTOR & PREFILLED SYRINGE



Even if you have given yourself an injection before, it is important to know that the Taltz autoinjector has different instructions from any device you may have used in the past.

**taltz**  
(ixekizumab)  
injection  
80 mg/mL

Take your device out of the refrigerator and let it sit for 30 minutes to warm to room temperature. Wash your hands, select an injection site, and clean it with an alcohol wipe. Let the injection site dry before you inject.

## AUTOINJECTOR

### 1 | UNCAP THE AUTOINJECTOR

Twist off cap and throw it away. Do not put the base cap back on, as this could damage the needle. Do not touch the needle.



### 2 | PLACE & UNLOCK

Place firmly at the injection site and unlock autoinjector.



### 3 | PRESS, HOLD, AND LISTEN

Press and hold the green button. Listen for a loud click. When you hear the second click, that means the full dose has been delivered.



**IMPORTANT:** These are not the complete instructions for using your device. Please see Instructions for Use that come with your device or visit [taltz.com](http://taltz.com). Before you or your caregiver inject for the first time, your healthcare provider should show you how to prepare and inject Taltz.

Once you've completed the injection, dispose of the used device in a Taltz sharps container or another puncture-resistant container with a secure lid. Do not dispose of Taltz in household trash. For more information on how to properly dispose of your Taltz device, visit [www.fda.gov/safesharpsdisposal](http://www.fda.gov/safesharpsdisposal).

## PREFILLED SYRINGE

### 1 | UNCAP THE SYRINGE

Do not put the needle cap back on, as this could damage the needle. Do not touch the needle.



### 2 | PINCH & INSERT

Pinch and hold a fold of skin, and insert needle at a 45-degree angle.



### 3 | PUSH IN THE PLUNGER

You will know your injection is complete when the green plunger rod shows through the syringe.



If you need a sharps container, you can get one free by calling Taltz Together at **1-844-TALTZ-NOW** (1-844-825-8966). For more information on Taltz Together, please turn to page 12.

## SELECT IMPORTANT STORAGE INFORMATION

Store Taltz in the refrigerator and protect from light. **Do not** freeze Taltz. **Do not** use Taltz if it has been frozen. **Do not** shake your Taltz device.


Please see the [Safety Summary](#) on pages 26 and 27. Please see [Instructions for Use](#) included with your device.

# DOSING FOR PSORIATIC ARTHRITIS

taltz  
(ixekizumab)  
injection  
80 mg/mL



Staying on your dosing schedule can help give you a chance at achieving the best possible results.



**GET STARTED DATE** \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

**MAINTAIN DATE** \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

If you have both psoriatic arthritis and moderate to severe plaque psoriasis, follow the dosing schedule for plaque psoriasis below.



For other important information about taking Taltz, talk to your doctor and refer to the Instructions for Use that come with your device. Before you or your caregiver inject for the first time, your healthcare provider should show you how to prepare and inject Taltz.

## SELECT IMPORTANT SAFETY INFORMATION


Your doctor should check you for tuberculosis (TB) before you start Taltz, and watch you closely for signs of TB during and after treatment with Taltz. If you have TB, or had it in the past, your doctor may treat you for it before you start Taltz.

Please see the [Safety Summary](#) on pages 26 and 27. Please see [Instructions for Use](#) included with your device.

# DOSING FOR ANKYLOSING SPONDYLITIS

taltz<sup>®</sup>  
(ixekizumab)  
injection  
80 mg/mL

Staying on your dosing schedule can help give you a chance at achieving the best possible results.



**GET STARTED DATE**

---

---

---

**MAINTAIN DATE**

---

---

---

For other important information about taking Taltz, talk to your doctor and refer to the Instructions for Use that come with your device. Before you or your caregiver inject for the first time, your healthcare provider should show you how to prepare and inject Taltz.

## SELECT IMPORTANT SAFETY INFORMATION

Your doctor should check you for tuberculosis (TB) before you start Taltz, and watch you closely for signs of TB during and after treatment with Taltz. If you have TB, or had it in the past, your doctor may treat you for it before you start Taltz.

Please see the [Safety Summary](#) on pages 26 and 27.  
Please see [Instructions for Use](#) included with your device.

## HELPFUL INJECTION TIPS

Talk to your doctor about whether the tips below may help with the injection of biologics.\*



Placing something cold (such as an ice gel pack or cold water bottle) on the injection site before and after the injection may help reduce pain.



Acetaminophen may help relieve pain.



Antihistamines or topical corticosteroids may help with itching or swelling caused by the injection.



Choose different injection sites each time you inject.

The recommended places for injection are your stomach (abdomen) or thigh, and another person may give you the injection in the back of your arm.

### TIPS SPECIFIC TO TALTZ



You should take Taltz out of the refrigerator and let it sit for 30 minutes to warm to room temperature before injecting. Leave the cap on the device while it is warming up to room temperature.



However, Taltz should not be microwaved, run under hot water, or left in direct sunlight.



There are 2 devices available for injecting Taltz: an autoinjector and a prefilled syringe.

\*These tips have not been developed or studied specifically for Taltz.

### SELECT IMPORTANT SAFETY INFORMATION

The most common side effects of Taltz include: injection site reactions, upper respiratory infections, nausea and fungal infections. Tell your doctor if you have any side effects. **You can report side effects at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

## THINGS TO KNOW

You may experience side effects while taking Taltz. If you do, please reach out to your doctor.

### INJECTION SITE REACTIONS

Injection site reactions may include injection site pain, redness, and possibly swelling of the skin.

- Injection site pain may occur while you are injecting. Other reactions, such as redness, may happen after several days
- Most injection site reactions did not cause people to stop taking Taltz
- In clinical trials, most injection site reactions were mild to moderate. Talk to your doctor about any injection site reactions

If you are experiencing injection site reactions with one type of device, talk to your doctor about whether it might be appropriate to switch to the other Taltz device.

### THE OTHER MOST COMMON SIDE EFFECTS INCLUDE:

- Upper respiratory infections
- Nausea
- Fungal infections

These are not all of the possible side effects of Taltz. Serious side effects may occur. Tell your healthcare provider about any side effect that bothers you or that does not go away. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [fda.gov/medwatch](http://fda.gov/medwatch) or call **1-800-FDA-1088**.

Lilly is currently monitoring and will continue to monitor and communicate additional information about injection site reactions. Please contact The Lilly Answers Center at **1-800-LillyRx** (1-800-545-5979) if you have more questions about this topic.

**taltz**<sup>®</sup>  
(ixekizumab)  
injection  
80 mg/mL

Please see the **Safety Summary** on pages 26 and 27. Please see **Instructions for Use** included with your device.


# TAKING TALTZ & STORAGE

taltz<sup>®</sup>  
(ixekizumab)  
injection  
80 mg/mL

Before using Taltz, be sure to read the Instructions for Use, visit [taltz.com](http://taltz.com) to watch the injection training videos, or have Taltz Together connect you to a nurse for an over-the-phone, in-person, or online discussion about injection training.

## WHEN TAKING TALTZ, PLEASE REMEMBER

- You may inject the medicine into your stomach (abdomen) or thigh
- Another person may give you the injection in the back of your arm
- **Alternate your injection site. Do not** inject in the exact same spot every time
- **Do not** inject within 1 inch of your belly button
- **Do not** give an injection into areas where the skin is tender, bruised, red, or hard, or in an area of skin that is affected by psoriasis
- Before you or your caregiver inject for the first time, your doctor should show you how to prepare and inject Taltz



## STORING TALTZ

- Keep Taltz stored in the refrigerator between 2°C to 8°C (36°F to 46°F) and make sure you do not freeze it
- When you are ready to use your device, take it out of the fridge and let it sit for 30 minutes to warm to room temperature before injecting
- If you drop your device on a hard surface, do not use it. Use a new one for your injection. Always handle carefully
- If needed, you may store Taltz at room temperature up to 30°C (86°F) for up to 5 days in the original carton to protect from light. Once Taltz has been stored at room temperature, do not return to the refrigerator and discard, if unused, within 5 days
- Keep Taltz and all medicines out of the reach of children



Do not use Taltz if it has been frozen



You should protect Taltz from light at all times



Do not shake your Taltz device

Remember to talk to your doctor about any other questions you may have.

## SELECT IMPORTANT SAFETY INFORMATION

Before you use Taltz, make sure your doctor knows if you are being treated for an infection, have an infection that does not go away or keeps coming back or have symptoms of an infection such as fever, cough, sores, diarrhea or other symptoms; have TB or have been in close contact with someone with TB; have Crohn's disease or ulcerative colitis.

Please see the **Safety Summary** on pages 26 and 27.  
Please see **Instructions for Use** included with your device.

Even if you have given yourself an injection before, it is important to know that the Taltz autoinjector has different instructions from any device you may have used in the past.

## PURPOSE AND SAFETY SUMMARY

**Important Facts About Taltz®** (tōl-ts). It is a prescription medicine also known as ixekizumab.

Taltz is an injectable medicine used to treat adults:

- With moderate to severe plaque psoriasis. It is for people who may benefit from taking injections or pills (systemic therapy) or treatment using ultraviolet or UV light (phototherapy).
- With active psoriatic arthritis.
- With active ankylosing spondylitis.

### Warnings

- Taltz affects the immune system. It may increase your risk of infections, which can be serious. Do not use Taltz if you have any symptoms of infection, unless your doctor tells you to. If you have a symptom after starting Taltz, call your doctor right away.
- Your doctor should check you for tuberculosis (TB) before you start Taltz, and watch you closely for signs of TB during and after treatment with Taltz.
- If you have TB, or had it in the past, your doctor may treat you for it before you start Taltz.
- Do not use Taltz if you have had a serious allergic reaction to ixekizumab or any other ingredient in Taltz, such as: swelling of your eyelids, lips, mouth, tongue or throat, trouble breathing, feeling faint, throat or chest tightness, or skin rash. Get emergency help right away if you have any of these reactions. See the Medication Guide that comes with Taltz for a list of ingredients.
- Crohn's disease or ulcerative colitis (inflammatory bowel disease) can start or get worse with Taltz use. Tell your doctor if you have any of these symptoms or if they get worse: stomach pain, diarrhea, and weight loss.
- You should not get live vaccines while taking Taltz. You should get the vaccines you need before you start Taltz.

### Common side effects

The most common side effects of Taltz include:

- Injection site reactions
- Upper respiratory infections
- Nausea
- Fungal infections

Tell your doctor if you have any side effects.

**You can report side effects at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

**taltz**<sup>®</sup>  
(ixekizumab)  
injection  
80 mg/mL

### Before using

#### Before you use Taltz, review these questions with your doctor:

- Are you being treated for an infection?
- Do you have an infection that does not go away or keeps coming back?
- Do you have TB or have you been in close contact with someone with TB?
- Do you have possible symptoms of an infection such as fever, cough, sores, diarrhea, or other symptoms? Ask your doctor about other possible symptoms
- Do you have Crohn's disease or ulcerative colitis?

#### Tell your doctor if:

- You need any vaccines or have had one recently.
- You are under age 18. Taltz has not been approved in people under 18 years of age.
- You take prescription or over-the-counter medicines, vitamins, or herbal supplements.
- You are pregnant or planning to become pregnant. It is not known if Taltz can harm an unborn baby.
- You are breastfeeding or planning to breastfeed. It is not known if Taltz passes into breastmilk.

### How to take

See the instructions for use that come with Taltz. There you will find information about how to store, prepare, and inject Taltz.

### Learn more

For more information, call 1-800-545-5979 or go to [taltz.com](http://taltz.com).

This summary provides basic information about Taltz and is not comprehensive. Read the information that comes with your prescription each time your prescription is filled. This information does not take the place of talking with your doctor. Be sure to talk to your doctor or other healthcare provider about Taltz and how to take it. Your doctor is the best person to help you decide if Taltz is right for you.

Taltz<sup>®</sup> is a registered trademark owned or licensed by Eli Lilly and Company, its subsidiaries and affiliates.

IX CON BS 02AUG2019

*Lilly*

taltz<sup>®</sup>  
(ixekizumab)  
injection  
80 mg/mL

Taltz<sup>®</sup> is a registered trademark owned or licensed by Eli Lilly and Company, its subsidiaries and affiliates.  
Taltz Together<sup>™</sup> and Companion in Care<sup>™</sup> are trademarks of Eli Lilly and Company.  
PP-IX-US-3825 04/2020 ©LILLY USA, LLC 2020. ALL RIGHTS RESERVED.

*Lilly*